

DNA and RNA-SIP reveal *Nitrospira* spp. as key drivers of nitrification in groundwater-fed biofilters

Arda Gülay^{1,4*}, Jane Fowler¹, Karolina Tatari¹, Bo Thamdrup³, Hans-Jørgen Albrechtsen¹, Waleed Abu Al-Soud², Søren J. Sørensen² and Barth F. Smets^{1*}

¹ Department of Environmental Engineering, Technical University of Denmark, Building 113, Miljøvej, 2800 Kgs Lyngby, Denmark. **Phone:** +45 45251600. **FAX:** +45 45932850. **e-mail:** argl@env.dtu.dk, jfow@env.dtu.dk, hana@env.dtu.dk, bfsm@env.dtu.dk*

² Department of Biology, University of Copenhagen, Universitetsparken 15, Building 1, 2100 Copenhagen, Denmark. **Phone:** +45 35323710. **FAX:** +45 35322128. **e-mail:** w.abualsoud@bio.ku.dk, sjs@bio.ku.dk

³ Nordic Center for Earth Evolution, Department of Biology, University of Southern Denmark, Campusvej 55, 5230 Odense, Denmark. **Phone:** +45 35323710. **FAX:** +45 35322128. **e-mail:** bot@biology.sdu.dk

⁴ Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, United States, 26 Oxford St, Cambridge, MA 02138, **Phone:** +1 (617)4951564. **e-mail:** ardagulay@fas.harvard.edu

Supplementary Figure 2




Fig.S2 Sequential work-flow to detect putative ammonium and nitrite oxidizers using RNA and DNA SIP. Detailed explanation of each filter step is given in Supplementary Materials and Methods.

